



Ethnobotanical Study of Medicinal Plants by Shenasha People Used to Treat Human Ailment in Dibati District, Northwest Ethiopia

Abesh Birhanu Morka^{1,*}, Tena Regassa Duressa^{2,*}

¹Ethiopia Biodiversity Institution, Mettu Biodiversity Center, Forest and Range Land Plant Biodiversity Case Team, Mettu, Ethiopia

²Department of Biology, Faculty of Natural and Computational Sciences, Wollega University, Nekemete, Ethiopia

Email address:

birhanuabesh12@gmail.com (A. B. Morka), tenaregasa@gmail.com (T. R. Duressa)

*Corresponding author

To cite this article:

Abesh Birhanu Morka, Tena Regassa Duressa. Ethnobotanical Study of Medicinal Plants by Shenasha People Used to Treat Human Ailment in Dibati District, Northwest Ethiopia. *American Journal of Environmental and Resource Economics*. Vol. 6, No. 3, 2021, pp. 91-102.

doi: 10.11648/j.ajere.20210603.13

Received: July 3, 2021; Accepted: July 26, 2021; Published: September 7, 2021

Abstract: *Background:* In all part of the world traditional medicinal plant practices formed the basis of health cares for human being before the advent of modern medicine. In the Ethiopia people use traditional medicine for treatment of human ailment. Like other communities in Ethiopia, shenasha people are use traditional medicinal plant in Dibati District, Benshangule gumuze Region, Ethiopia. The objective of the study was to document traditional knowledge on medicinal plants in Dibati district northwest. *Methods:* The main data collection tools were semi structured interviews, discussions, and observation focused group discussion and various ranking and comparison methods were employed. Information was collected from a sample of 99 informants (77 males and 22 Females) included 20 (male 16 female 4) key informants are identified by the help of kebele administration. *Results:* Among the study of analysis 48 medicinal plant species with 46 genera and 28 families were recorded. In terms of this the Most of the species (36) were collected from the wild while 12 were collected from home garden were collected. Among the most widely used method of preparation was crashed (33.33%), Pounding (18.75%), of the different medicinal plant parts. *Conclusion:* In the study area possesses diverse natural vegetation and the environment under serious threatened by mainly agricultural expansion, firewood collection, population pressure, overgrazing, urbanization, household tool construction, charcoal production and medicinal purpose. Awareness rising on the use and management systems, sustainable utilization of medicinal plants and their in-situ and ex-situ conservation and establishment of forest protected areas should be recommended.

Keywords: Medicinal Plant, Dibati Woreda, Informant, Shinasha People, Indigenes Knowledge, Ethnobotanical

1. Introduction

In all part of the world traditional medicinal plant practices formed the basis of health cares for both human being and animal before the advent of modern medicine. In the Ethiopia people have been using traditional medicine to treat both human and animal disease for generations. Traditional medicine is still wildly practices in rural area where modern medicine and services available [27]. Many of today's wonderful drugs were initially discovered through the study of traditional medicine [27]. Availability of medicinal plant has been affected by a dramatic decrease of native vegetation

due to agricultural explanation, deforestation, fire, and overgrazing drought, tread of charcoal, firewood, introduction of alien invasive species and urban associated development. Medicinal plants obtained from wild habitats are found in different natural ecosystems of the forests, grasslands, woodlands, wetlands, in field margins and garden fences, as weeds and in many other microhabitats from where they are harvested when the need arise. There are many medicinal plants of Ethiopia that have good properties for land rehabilitation and erosion control Maintain health through traditional medicine in general and utilization of medicinal plant in particular is almost as old as the history of

man kinds [9]. This is true in Ethiopia 80% of the population still relies on the plant to prevent and cure various health problem [15]. The current plant use trend in Dibati district shows that the environment is facing problems of resource depletion and loss of indigenous knowledge like other areas of the country. Thus, intensive ethnobotanical research plays a vital role to draw information on plants and related indigenous knowledge for conservation and sustainable utilization. Like many other parts of the country, there is no such ethnomedicinal research and documentation carried out in Dibati District, Metekele Zone, Northwest and Benshangul Gumuz Region. Dibati district is one of our research areas which are a very high amount of Shinasha people lives. In this woreda there is no any research which is done on the use of medicinal plants to treat human ailment by the Shinasha people's practices.

2. Methods

2.1. Description of the Study Area

The Benishangul Gumuz Regional State (BGRS) was established in 1994 as one of the nine regional states of Ethiopia. The BGRS borders the Republic of Sudan in the west, Amhara region in the North, Oromiya in the South east and Gambella region in the South. Administratively, it is divided in to 3 zones (Metekele, Assosa, and Kemashi) that are divided in to 19 Woreda, and one special Woreda. Metekel Zone is divided into 7 Woreda out of which Dibati Woreda, the study area is located North 100 55'-110 90' latitude North and 300 12'-300 36' longitude East.(Source BGRS, 2008).




Figure 1. Map of the study area.

2.2. Reconnaissance Survey and Selection of Study Sites

Dibati Woreda has a total of 29 kebeles. Out of these, 5 kebeles were selected those are Modorem, Gerze, Zegehe, Legbuna, and Bechati due to the have high amount of shinasha peoples found in kebele. Additional to that they have larger vegetation cover and presence of key informants, different altitudinal ranges are selects for ethnobotanical data collection. The traditional healer, used as key informants, was identified by with the assistance of local authorities, elders and knowledgeable persons.

2.3. Informant Selection

Information regarding to each peasant association leaders

the total household of Shenasha people in the study area of five kebele from the age of twenty and above is (N) is male 651 female 141 total 792. To obtain total sample size from the total targeted household, the researcher was use the usual known formula [30] that was calculated as follows:

$$n = \frac{N}{1 + N(e)^2}$$

Where n is sample size of the study area

N is targeted population size (total population of five kebele in study area.)

e: is the level of precision or sampling error= (0.05)

For the above formula,

$$n = \frac{792}{1 + 792(0.05)^2} = 99$$

A total 99 informants five to seven individuals from each study kebele from the age of twenty and above were included. From the five studies kebele twenty key informants were purposively selected based on recommendations from local authorities (kebele administrators and local guides) peasant association leaders and other members of the local communities.

Plant specimen collection and identification

At the end of the interview, the reported medicinal plants were collected from natural vegetation and home garden. Sample specimens of the plants cited for their medicinal use was collected, numbered, pressed and dried for identification. Plant identification was performed both in the field, and at the National Herbarium of AAU. Preliminary identification was done in the field and reconfirmed at the National Herbarium. Identification of other plant specimens with my advisor Dr. Tena Regass and Wollega University herbarium workers was done in November 2017, using the Flora of Ethiopia and Eritrea and also by comparison with authenticated specimens.

2.4. Ethnobotanical Data Collection

Ethnobotanical data was collected between October to November, 2017 on three field trips. The data was collected based on prepared questionnaires, semi-structured interviewees, observation, focus group discussion, and guided field walks with informants were employed to obtain indigenous knowledge of the local community of shinasha people. The study was carried out by interview in their habitations in different village. The informants include the various data sets such as local names, disease treated, parts of the plant used, and method of preparation dose and route of application was obtained from local people through individual interviews. A list of question was prepared that covers the discussion with the informants in particular orders. All of the interviews were held in Amharic and Shinasha language of the local people. The place and time for discussion were set based on the interest of the informants. [7].

2.5. Ethnobotanical Data Analysis

The Ethnobotanical data collected was analyzed following survey and analytical tools for ethnobotanical methods which are Informant's preference ranking, descriptive statistic (Microsoft excel spreadsheets software), Direct matrix ranking and paired comparisons conducted following [19, 7, 22]. Were computed to assess the degree of effectiveness of certain medicinal plants against human. Additional to that the use value (UV), a quantitative method that demonstrates the relative importance of species known locally [17]. was calculated using the following formula. $UV = \sum U/n$ where: UV = use value of a species; U = number of citations per species; n = number of informants.

Fidelity level index quantify the importance of a given species for a particular purpose in a given cultural group [12, 7].

The formula for FL is given as $FL\% = Ip / Iu \times 100$, where Ip the number of informants who independently indicated the use of a species for the same major ailments and Iu the total number of informants who mentioned the plant for any major ailment [12].

Informant consensus factor (ICF) was calculated for categories of ailments to identify the agreements of the informants on the reported cures using the formula used by [17, 19]. ICF was calculated as follows: number of use citations for each ailment (nur) minus the number of species used (nt) for that ailment, divided by the number of use citations for each ailment minus one. $ICF = (nur - nt) / (nur - 1)$ where: ICF: Informant consensus Factor, Nur: number of use citation Nt: number of species used.

3. Results and Discussion

3.1. Medicinal Plant Knowledge Among Informants

In study area comparison of educational status, non-educated informants handled much knowledge of traditional medicine whereas educated informants had low knowledge of traditional medicine, which is an indicative of impact of modern education. Similar findings were reported by [28, 26, 14, and 15].

Regarding to sex male are more knowledgeable medicinal plants than females because of cultural influence and they can do more home activity. Similar results were reported by [14, 24].

Medicinal plants used to treat human ailments in the Dibati Woreda

The 48 of ethnomedicinally important plant species that are used to treat 57 human ailments recorded in five kebeles of Dibati woreda. These plants belong to 46 genera and 28 families. Family Fabaceae contributed 7 species, Astraceae 3 species, Combretaceae 3 species, Euphorbiaceae Bignoniaceae species, Amaranthaceae, Rubiaceae, Alliaceae, Cucurbitaceae, Olaceaceae, Lamiaceae, Polygonaceae, Ranunculaceae, Rutaceae and Solanaceae comprises 2 species in each, and represented by 2 species each and the rest 15 families comprises 1 species each. Some of the medicinal plants recorded are also used as remedies in other parts of Ethiopia. [11], documented 30 plant species, [20], 78 species and 12 plant species are mentioned [13].

The dominance of family Fabaceae for the treatment of human diseases was reported in the work of [8, 3], and others [20]. This might be due to their weedy nature of the family Fabaceae and take advantages of disturbance.

3.2. Major Human Ailments and Plant Species Used by Local People in the District

Though more than 57 different diseases of humans were recorded as human health problem that are treated by 48 plant species, one species can treat a single disease or a number of diseases. The practitioners of the area commonly diagnose each health problem by an interview and visual inspection of the patient. These shows large numbers of

diseases have got solution by traditional medicine in Dibati Woreda compared to different investigations in Ethiopia. For example, [14] reported 135 plant species that used to treat 65 ailments of humans and [24] reported 80 plant species that used to treat 49 ailments of humans.

According to the informants the largest number of species was used to treat snake biting, which is treated with 11 species. This disease is common in the area because most people engage in living the farmland as a result their men are susceptible to the snake biting; the next is eye vile is treated by 9 species. Amoebae, toothache, febrile illness and wound each disease are treated by 7 species and malaria, stomach ache common cold, retained placenta and abdominal pain each disease are treated by 5 species.

In addition to these, the practitioners were also visited more for diseases like gastritis, eye problem, homerriod, athlete foot and others.

3.3. Habitat, Habit, Plant Part Used, Methods of Preparation, Rout of Administration and Application of Human Medicinal Plants in the Study Area

From the medicinal plants that are used for human ailments, 36 species were collected from the wild vegetation and 12 species from home gardens. These indicated that the local people obtain medicinal plant species from wild vegetation than home gardens. This result agrees with [10, 8, 28, 26, 14, 24]. This indicated that the practice of cultivation of medicine plants for their medicinal purpose in home gardens of most of the country is low although many plants are cultivated for other purposes, mainly for food. In a similar way people in the study area have less effort to cultivate medicinal plants in their home gardens rather go to the nearby or far places and harvest the plants.

3.4. Habit Human Medicinal Plants

In the study area there are many habits of medicinal plant those are herb, shrub, tree and climber. The result shows that analysis of growth forms of medicinal plants revealed that herbs constitute the largest category 19 (39.58%) followed by tree 11 (22.92%), 12 (25.00%) shrub and 6 (12.50%) Climbers were recorded. Some researchers previously studies [5-10, 4, 13, 25, 21, 14, 24, 6].




Figure 2. Habit of medicinal plants for human in the Dibati woreda.

3.5. Plant Part Used to Treat Human Ailments

From the total plant parts used for remedy preparation the leaves and the roots were the most commonly used plant parts in the preparation of remedies. The most widely used plant part for the preparations of remedy were leaves, which accounted for 10 (20.85%) followed by roots 8 (16.66%), seed 6 (12.50%) break 4 (8.33%) bulb 4 (8.33) and others. The most part were leafe because of the availibility of faild collection, ease to preparation and the avialibilty of medicinal content. This result agrees with following researcher [5, 10, 25, 8, 4, 13, 21, 14, 24, 6].




Figure 3. Percentage of medicinal plant parts used for human health treatment.

3.6. Method of Preparation

The preparations vary based on the type of disease treated and the actual site of the ailment. The principal methods of

plant parts remedy preparation forms were reported to be through crushed, which accounts for 16 (33.33%) followed by pounded 9 (18.75%), squeezing 7 (14.58%), chewing 5 (10.42%), crashing and pounded 4 (8.33%), cocking 3 (6.25%),

eating 1 (2.08%), smoke/ burning 1 (2.08%) crushed squeezed 1 (2.08%), pounded and squeezed 1 (2.08%) and pounding powder 1 (2.08%) respectively. The most domina method of preparation is crushed. The result showed that the majority of remedies were prepared from single plant species and few are

prepared from different, Solvents and additives which is a combination of medicinal plants, was used to treat a disease. In agreement with this study, similar researcher of medicinal plant species method of preparation is reported from different parts of the country by [14, 24, 6, 31].




Figure 4. Method of preparation of medicinal plant used for human health treatment.

3.7. Rout of Administration

There are various routes of administration of traditional medicinal plants prepared products by the local community. The major routes of administration in the study area are oral, dermal, nasal, anal, tide; ear and fumigated People of the study area

mostly administer traditional medicine orally. Oral accounts 25 (52.08%) followed by dermal 6 (16.67%), fumigated 4 (8.33 oral and dermal 2 (4.17%) and others (Figure 8). This fact that has been documented by different authors in the other part of Ethiopia [4, 13, 2, 1, 26, 21, 6, 23, 24].




Figure 5. Rout of administration of medicinal plant used for human health treatment.

3.8. Dosage of Medicinal Plants Used in Human Ailments

People of the study area used various units of measurement and the duration of administration to determine the dosage. Local units such as finger length (e.g., for bark, root, stem), different measuring materials (e.g., spoon, coffee cup, tea cup and glass cups) and numbers (e.g., for leaves, seeds, fruits, bulbs were used to estimate and fix the amount of medicine. But, these measurements are not accurate enough to determine

the precise amount. For medicinal plants that are taken topically they do not have clear cut dosage [20, 29, 1, 28, 14], we have also discussed lack of precision and standardization as one drawback for the recognition of the traditional healthcare system. Although the most of remedies were reported to have no serious adverse effect vomiting and temporary inflammations. These could be attributed to low toxicity of the remedies preparation of medicinal plant species used by traditional healer in the study area. However, the toxicity of

some medicinal plant and the potential to do harm is a common complaint among those who would like traditional medicine to be standardized. It is commonly believed that traditional practitioner either does not know the strength of their own medicine or does not bother to fit doses to the size or the body weight of the patient. However it is that some traditional healers do give different dosage and frequencies of application depending on age sex pregnancy and other condition or the medicinal itself on such difference. However, from the interview made during the study, it was found that there was disagreement among the healers concerning the dosage system used. For example, some informants suggested that two or three glass of the squeezed from *Justicia schimperiana* is used to treat, snake bit, malaria, stabbing pain and toxic substance while some suggested that only one glass is enough for the same problem. Still some others suggested that they apply the glass randomly without such measuring system or without mentioning a fixed by saying “squeezed of glass”. Although the measurements used to determine the dosages are not standardized and doses given depend on the age, physical appearances and health conditions; that is, children are given less dose than adults, physically strong individuals take more dose than weak individuals depending on the type of disease. Though such prescription difference was practiced, still the amount prescribed by healers for both children and adults might not conform to the standard prescriptions as in modern medical literature. The absence of any adverse effects of traditional medicines after administration were also more frequently mentioned by the traditional healers but some of the preparations were reported to have some adverse effects like diarrhea, headic, abdominal pain and vomiting. The

traditional healers indicated that they use antidotes for the adverse effects of some traditional medicines like eating cooked teff flour and honey, drinking boiled coffee, “yedoro weat” milke <ergo> suger after taking the medicine. For instance, the use of *Euphorbia abisanica* for the treatment of gonorrhea, the sanp is collected mixed with “teff” powdered and backed and then eaten before any food for 3 days. During that diarrhea follows, as an antidote the local healers ask the patient to drunk boiled coffee.

In addition to the use of *Jatropha curcas* for the treatment of Rabbis, one seed are pounded, powdered, mixed with milk then the filtrate the solution is drunk. During that dihareha follows, as an antidote the local healers ask the patient to drunk borede and milk. In addition to the use of *Nicotino tabacum* for the treatment of snake bit, one leaf is pounded, powdered, mixed with water then the filtrate the solution is drunk. During that vomiting follows, as an antidote the local healers ask the patient to drink milk.

3.9. Application

The prepared traditional medicines are applied in a number of methods, drinking account the largest that account 12 (25.00%), followed by smoking 10 (20.83%), sniffing 7 (14.58%), dropping 5 (10.42%), painting 4 (8.33) swallowing 3 (6.25) eating 2 (4.17) put on 1 (2.08) tide 1 (2.08) washing accounts 2 (4.17) burshing 1 (2.08)% (seen Figure 6). Internal ailments were commonly treated by making the patient drink herbal preparations; tooth infection were treated by crushing and put on the remedial plant part on the tooth surface; skin infections such as ringworm were treated by painting herbal preparations on an infected skin.




Figure 6. Application of medicinal plant used for human health treatment.

Sources and Transferring of Indigenous knowledge Sources of Traditional knowledge practices.

The highest number of traditional medicinal plant knowledge gain from to be 45 (45.4%) Father followed by 20 (20.2%) from Mother and 15 (15.1%) from Uncle 10 (10.1%) from Brother, 8 (8.08%) from Sister 6 (6.06) from Neighborhood and 5 (5.05%) from Friend. The great majority of respondents (90%) reported that most of their knowledge was received from their family members and friends secretly by oral. The secret practices of traditional medicines came from their ancestors. Within my finding similar studies are agreed by [28, 26, 31, 24,].




Figure 7. Sources of knowledge on the practice of traditional medicine.




Figure 8. Transferring of knowledge of traditional medicinal plants.

3.10. Transferring Knowledge of Traditional Medicinal Plants

According to the survey, knowledge transfer of medicinal plants follows vertical transfer to the most selected family member orally with great secrecy. The findings of the study showed that as people become older and older their knowledge of traditional medicine becomes better and better. Most of the informants were elders that indicated the trend of transferring knowledge is usually at old age. The highest number for the ways of transferring knowledge on traditional medicinal plants by elder son who received 56 (56.5%) votes followed by the elder daughter 12 (12.1%), for the brother 10 (10.1%) for the sister 7 (7.07%) for the not to all 5 (5.05%) for all went to known 4 (4.04%) to all the member of my family 3 (3.03%) and other to fried 2

(2.02%) (Figure 7). Therefore most way of indigenous knowledge transfer in the study District was by word of mouth to a family member (especially to an elder son). Similar findings were reported for other communities in Ethiopia [26, 24, 31].

4. Ranking of Most Important Medicinal Plants

4.1. Preference Ranking

Preference ranking for eleven medicinal plants to treat snake biting (Table 6) made by ten informants showed that *Bersema abyssinica* ranked first and hence is the most effective medicinal plant to cure snake biting. *Myrsine melanophloeos*, *Nicotiana tabacum*, *Justicia schimperiana* and *Allium sativum*. Are the 2nd, 3rd, 4th and 5th respectively.

4.2. Direct Matrix Ranking

Direct matrix ranking was performed to assess the relative importance each of the plant. The result of the direct matrix ranking showed that *Cordia africana* stood first in being the most multipurpose medicinal plant followed by *Gardenia ternifolia*, *Eucalyptus globulus*, *Myrsine melanophloeos*, *Croton macrostachyus* Del. *Acacia abyssinica*, *Ximenia caffra. sond*, *Bersema abyssinica* and *Jatropha curcas* was the least (Table 7).

Table 1. Preference ranking of eleven Selected Medicinal Plants based on the Degree of their Curative Power of snake bit as Perceived by Informant.

Name of plants species	Respondents (R1-R10)										Total	Rank
	R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	R ₉	R ₁₀		
<i>Allium sativum</i> L.	4	3	5	3	4	5	1	5	5	3	38	5 th
<i>Carduus englarium</i> Sch. Bip. Ex A. Rich.	2	3	1	1	2	3	4	3	3	5	27	9 th
<i>Carissa spinarum</i> L.	2	5	5	4	5	5	2	5	3	1	37	6 th
<i>Justicia schimperiana</i> (Hochst. ex Nees) T. Anders.	4	5	5	2	5	4	5	4	2	3	39	4 th
<i>Lagenaria siceraria</i> (Mol.) Standl.	4	4	0	1	4	3	2	2	1	3	24	11 th
<i>Nicotiana tabacum</i> L.	3	5	2	4	4	5	5	3	5	5	41	3 rd
<i>Paveonia urens</i> cav	5	2	5	3	2	4	5	4	1	5	36	6 th
<i>Stereospermum kunthianum</i>	4	2	2	4	5	2	2	5	5	4	30	9 th
<i>Bersema abyssinica</i>	5	5	5	4	5	5	4	4	5	5	47	1 st
<i>Dicrocephale latifolia</i>	5	4	2	2	3	2	3	3	3	4	31	8 th
<i>Myrsine melanophloeos</i>	5	4	4	5	5	5	4	5	5	4	46	2 nd

Table 2. Direct matrix ranking for seven specie and main use in study area.

NamePlant species	Use categories							Total	Rank
	Charcoal	Construction	Medicine	Food	Firewood	Fencing	Furniture		
<i>Jatropha curcas</i>	0	0	4	0	3	5	1	13	9 th
<i>Ximenia caffra. sond</i>	0	2	5	5	5	3	2	22	7 th
<i>Cordia africana</i> Lam.	5	5	5	4	5	3	5	32	1 st
<i>Croton macrostachyus</i> Del.	5	5	5	0	4	4	2	24	5 th
<i>Acacia abyssinica</i>	5	5	3	0	4	3	3	23	6 th
<i>Eucalyptus globulus</i>	5	5	3	0	5	5	5	28	3 rd
<i>Myrsine melanophloeos</i>	5	5	5	0	5	2	3	25	4 th
<i>Bersema abyssinica</i>	1	2	5	0	3	3	1	15	8 th
<i>Gardenia ternifolia</i>	2	3	5	3	4	3	5	30	2 nd
Total	33	30	40	12	47	31	23		
Rank	3 rd	5 th	2 nd	7 th	1 st	4 th	6 th		

4.3. Paired Comparison

A paired comparison made to determine the most preferred medicinal plants among the 9 species that were used to treat eye vile in the study area, the responses of ten key informants, showed that Ranked *Withania*

somnifera first followed by *Gardenia ternifolia*. (Table 3). Therefore, this result indicated that *Withania somnifera*. is the most preferred while. *Clausena anisata* is the least favored over the other plant species cited in treating eye vile.

Table 3. Paired comparisons of five medicinal plant species used to treat fibril illness.

Name of plants species	Respondents (R1-R10)										Total	Rank
	R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	R ₉	R ₁₀		
<i>Gardenia ternifolia</i>	1	4	3	0	1	1	2	4	5	4	35	2 nd
<i>Acacia abyssinica</i>	4	1	2	2	4	5	5	4	2	1	30	4 th
<i>Allium sativum</i> L.	2	2	1	4	3	2	4	3	2	5	28	5 th
<i>Croton macrostachyus</i> Del	5	5	1	4	2	2	1	0	1	4	25	8 th
<i>Vernonia amygdalina</i> Del.	1	0	2	4	3	1	4	1	5	1	27	6 th
<i>Secuidaca longepeduncula</i>	4	1	0	4	1	2	1	4	5	4	26	7 th
<i>Clausena anisata</i>	4	4	1	0	4	1	4	3	1	0	22	9 th
<i>Capparis tomentosa</i>	1	2	4	2	4	2	4	4	3	5	31	3 rd
<i>Withania somnifera</i>	4	2	4	4	5	1	5	4	4	5	38	1 st

5. Threatened and Factor Threatening Medicinal Plants in Dibati Woreda

5.1. Threatened Medicinal Plant in the Study Area

The ranking of 9 medicinal plants based on the degree of threats was conducted using 10 key informants. The results (Table 4) indicated that is *Asparagus africanus* and *Myrsine melanophloeos* the most threatened followed by *Withania somnifera* (L.) and *Bersema abyssinica* and the least threatened one is *Glinus lotoides* L.

Table 4. Ranking of threatened plants.

Name of plants species	Respondents (R1-R10)										Total	Rank
	R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	R ₉	R ₁₀		
<i>Myrsine melanophloeos</i>	1	4	3	0	1	1	2	4	5	4	35	2 nd
<i>Bersema abyssinica</i>	4	1	2	2	4	5	5	4	2	1	30	4 th
<i>Paveonia urens</i> cav	2	2	1	4	3	2	4	3	2	5	28	5 th
<i>Oncoba spinosa</i> Forssk.	5	5	1	4	2	2	1	0	1	4	25	8 th
<i>Momordica foetida</i> . schum	1	0	2	4	3	1	4	1	5	1	27	6 th
<i>Kalancheo petiti</i> A. Rich,	4	1	0	4	1	2	1	4	5	4	26	7 th
<i>Glinus lotoides</i> L.	4	4	1	0	4	1	4	3	1	0	22	9 th
<i>Withania somnifera</i> (L.)	1	2	4	2	4	2	4	4	3	5	31	3 rd
<i>Asparagus africanus</i>	4	2	4	4	5	1	5	4	4	5	38	1 st

5.2. Factors Threatening Medicinal Plants in the Study Area

The major reported threats of medicinal plants in the current study areas are agricultural expansion, including clearance of the major habitats of the species, and low attention of the local communities in conserving and protecting the species. However, there were reportedly indirect conservation activities of the species. These included the planting of the species for indirect uses such as aesthetics, fence, food, shade, spice, fuel wood, and source of income. Informants ranked agricultural expansion, fire wood and population pressure as the most

serious threat to the medicinal plants followed by medicinal purpose and charcoal collection is lower levels of threats by the other factors (Table 5). Similar study [32, 16, 2, 26, 31]. This showed that, there are different threats in medicinal plants such as agricultural expansion fire wood collection and others.

The agricultural expansion and deforestation was the major medicinal plant treats. The finding was in line with other findings [19]. This might be due to continuous agricultural expansions, deforestation and draught in addition to lack attention towards the medicinal plants. The plants are disappeared because of rapid socioeconomic, environmental and technological changes and as a result of the loss of cultural heritage under the guise of civilization.

Table 5. Ranking of threats to medicinal plants.

Name of plants species	Respondents (R1-R10)										Total	Rank
	R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	R ₉	R ₁₀		
Fire wood	4	5	5	4	3	4	3	5	2	5	40	2 nd
Medicinal purpose	2	1	2	1	2	2	1	2	2	1	16	8 th
Charcoal Making	4	2	3	2	3	1	3	4	1	3	23	7 th
Household tool construction	4	5	5	3	1	2	4	4	2	3	33	6 th
Over grazing	3	4	4	2	5	5	4	3	2	3	35	4 th
Agriculture expansion	5	4	5	5	5	4	4	5	5	5	47	1 st
Urbanization	3	2	4	5	4	3	2	2	5	4	34	5 th
population pressure	5	2	3	4	5	5	3	3	4	5	39	3 rd

5.3. Management and Conservation of Medicinal Plants

As a result many medicinal plants are under serious threats. So the local people should be conserve medicinal plant in-suit and ex- suit management style. Those manes that forty eight medicinal plants that are collected in the wild by conserve in-suit conservation method and the reaming seventeen medicinal plants that conserve by ex-suit conservation method. According to the informant information generally, there are some conservation measures that have been under taken around the world aimed at protecting threatened medicinal plant species from further destruction by create awareness for the user local people for the use and management of medicinal plant in study area.

Some authors clarify that home gardens can be refuge for wild species that are threatened in the wild by deforestation and environmental changes. Concerning this [33]. Reported that home gardens are being used as informal experimentation plots for new varieties and exotic species. Informants also reported that the healers know time and processes of gathering, and storing medicinal plants. It is once a year that some medicinal plants are collected and preserved. *Lepidium sativum*, *Cucuribita pepo*, *Jatropha curcas* and *Ocimum basilicum* seed, leaf, fruit or root are harvested, dried and preserved in roof corners or outside house, and dried parts are powdered and stored in different containers like pots, bottles or tied with clothes and used when needed. The study indicated that many of the informants who have knowledge on traditional medicine usage give priority to the immediate use of the medicinal plants than to its sustainable future uses, as a result their harvesting style is destructive. However, some plants has

protected for their spiritual and cultural purposes. Thus, these places are good sites for the protection of the medicinal plants since cutting and harvesting are not allowed in such particular areas. This was indicated that a good practice for the conservation of medicinal plants through cultivation [26].

6. Informant Consensus Factor (ICF)

All cited human and livestock diseases were categorized into 7 categories: namely, Sense organs related diseases, Animal and insect biting related disease, digestive system related disease, Reproductive system related diseases, Cultural related disease, Respiratory disease related diseases, intestinal and parasitic infection related diseases, and derma related disease, These diseases are categorized based on nature of disease, conditions that cause, place of attack, symptoms and sign of disease (Table 6). Disease categories with relatively higher ICF values were: intestinal and parasitic infection related diseases (0.89), derma related diseases (0.87) Sense organs related diseases (0.83), Cultural related diseases (0.73). This may indicate the common occurrence of these diseases so that more number of people exchanging information and agree on plant species that can be used to treat these diseases than the rest. The medicinal plants that are presumed to be effective in treating a certain disease have higher ICF values. On the other hand, the rest of disease categories had ICF value of lower suggesting that these diseases are either rare in the study area or are treated only by the healers with little information passed to other general public.

Table 6. Informant consensus factor (ICF).

Categories	Ailments/diseases	Number of Species used (nt)	Use citation (nur)	ICF Value
Sense organs	Eye problem, ear problem, trachoma,	5	25	0.83
Animal and insect biting	snake bite, rabies, malaria, spider poison, scorpion poisons	23	69	0.64
digstive system	Stomach ache,, diarrhea, toothache, gastrit abdominal pain	25	81	0.70
Reproductive system	gonorrhea, sexual weekens diseases, retained placenta, infertility RH factor	11	33	0.68
Respiratory diseases	Nasal bleeding,, tonsillitis, cough and asthma, common cold, leech	16	28	0.44
intestinal and parasitic infection	Tap worm, ascarisis	6	49	0.89
dermal	Dandruff, wound, skin rash, body swell, bone broken Leprosy goiter, ring worm mouth rash Athletes foot	8	58	0.87
Cultural releted	Eye vile, feberal ilnes, headache, preventing snake	18	64	0.73

6.1. Fidelity Level Index (FLI)

Fidelity level (FL) quantifies the importance of a species for a given purpose. Hence, fidelity level values were calculated for commonly used individual medicinal plants against the following ailments: *Justicia schimperiana* (against snake biting and Rabbis) *Asparagus africanus* (against retained of placenta) *Croton macrostachyus* (against

eye vile), *Stereospermum kunthianum* (Tonsil, scorpion biting), *Momordica foetida. schum* (against Bleeding during delivery) *Euphorbia abyssinica* (against Hemorrhoid, gonorrhea) *Paveonia urens* (against Rheumatic, tooth ache) and *Protulaca sp.* (against Gastritis). In the fact that these medicinal plants had the highest FL values which could be an indication of their good healing potential (Table 7).

Table 7. The relative healing potential of individual medicinal plants used against human or livestock ailments.

No	Medicinal plants species	Aliments treated	IP	IU	FLI (%)	Rank
1	<i>Croton macrostachyus</i>	Evile eye	52	58	89.6	3 th
2	<i>Justicia schimperiana</i> (Hochst. ex Nees) T. Anders.	Snake biting, rabbis	47	47	100	1 st
3	<i>Asparagus africanus</i> (kunth) Baker	Retained placenta	38	41	92.6	2 rd
4	<i>Stereospermum kunthianum</i> Cham.	Tonsil, scorpion biting	31	35	88.57	4 th
5	<i>Euphorbia abyssinica</i> J. F Gmel	Hemorrhoid, gonorrhea	29	37	78.38	6 th
6	<i>Momordica foetida. schum</i>	Bleeding during delivery	26	31	83.8	5 th
7	<i>Protulaca sp.</i>	Gastritis	21	29	72.4	7 th
8	<i>Paveonia urens</i>	Rheumatic, tooth ache	19	25	76.0	8 th

6.2. Medicinal Use Value

While some plant species are known to treat a single ailment, some others may be used for multiple of health problems. Medicinal use value is a quantitative method that demonstrates the relative importance of species known locally [33]. Some species that were cited for more than one ailment were selected and their use value was calculated. Results of use value computation for these species showed that *Croton macrostachyus* and had the *Carissa spinarum* highest use value (Table 8). This suggests that these species

are used to treat many ailments. For example, *Croton macrostachyus* was reported to treat ailments such amoeba, evil eye, febrile illness, wound, prevent snake, skin infection and malaria. On the other hand was *Carissa spinarum* L. reported to treat snake biting, ascarises, malaria, gonorrhea, Amoeba, febrile illness and diarrhea. *Justicia schimperiana* was reported to treat ailments such as snak biting, anti-toxic stabbing malaria, rabbis and typhoid's. *Stereospermum kunthianum* was reported to treat ailments such as scorpion biting, retained plasta, gasterite, tostile, snake biting and stomachache.

Table 8. Use Value of Certain Medicinal Plants in Study Area.

Plant species	Use citation (U)	Number of informants (n)	Use value (Uv)
<i>Croton macrostachyus</i>	7	23	0.35
<i>Carissa spinarum</i> L.	7	19	0.37
<i>Stereospermum kunthianum</i> Cham.	6	12	0.50
<i>Justicia schimperiana</i> (Hochst. ex Nees) T. Anders	6	14	0.43
<i>Vernonia amygdalina</i> Del.	5	12	0.42
<i>Combretum collinum</i>	5	13	0.38
<i>Paveonia urens</i>	5	15	0.33
<i>Cordia africana</i> Lam.	5	23	0.22
<i>Solanu nigrum</i>	4	11	0.36
<i>Kalanchoe peltata</i> A. Rich	4	9	0.44
<i>Gardenia lutea</i> Fresen.	4	12	0.33
<i>Allium sativum</i> L.	3	7	0.43
<i>Clusia abyssinica</i> Jaub. and Spach	3	5	0.60
<i>Coffea arabica</i> L.	3	25	0.12
<i>Ximenia caffra</i> Sond.	3	8	0.38
<i>Nicotiana glauca</i> L.	3	7	0.43
<i>Ruta chalepensis</i> L.	3	19	0.16
<i>Eucalyptus globulus</i> Labil.	3	20	0.15
<i>Zingiber officinale</i> Roscoe,	3	15	0.20
<i>Clematis integrifolia</i> Perr. and Guill	3	12	0.25
<i>Euphorbia abyssinica</i> J. F Gmel	3	4	0.75
<i>Linum usitatissimum</i> L.	3	11	0.27
<i>Oncoba spinosa</i> Forssk.	3	9	0.33

7. Conclusion

A study on medicinal plant utilization in area revealed that the shinasha communities commonly use medicinal plants for maintaining their primary healthcare. The study resulted in documenting 57 medicinal plants species 46 genera 28 family where Fabaceae is the leading family with the highest proportion of medicinal plants. The majorities of these medicinal plant species were obtained and collected 36 from wild, 12 from home garden. Analysis of growth forms of these medicinal plants that herbs constitute the largest category 19 (39.5%) followed by tree 12 (25.00%) shrub 16 (24.61%) plant species. Herbal remedies are prepared from fresh materials 45 (62.23%) and dried plant materials 8 (12.30%) and in both condation 18 (18.46%). In the study area, 72 ailments were reported (57 for human) which are being treated by traditional medicinal plants of the area. Leaves were the most frequently used plant parts followed by roots for preparation of human and livestock remedies. Most of the medicinal plants are administered orally (52.08%) and followed dermal (1667). The most widely used method of preparation was crashed (33.33%), Pounding (18.75%), of the different medicinal plant parts. They have a wide knowledge in using plants for various purposes such as for medicine, food, household utensils, fodder, fuel, construction, etc. This knowledge is transferred from elders to youngsters entirely through oral traditions and personal experiences. The major threats to medicinal plants and the associated knowledge in the study area are mainly agricultural expansion, firewood collection, population pressure, over gaze ring, urbanization, household tool construction, charcoal production and medicinal purpose. Therefore, use and management system awareness rising should be made among the healers so as to avoid erosion of the indigenous knowledge and to ensure its sustainable use.

Conflicts of Interest

The authors declare that they have no competing interests.

Acknowledgements

We would like to express our sincere appreciation to dibati worda administration office for the financial support during this study. We are thankful to the local communities of the district, Meteke Zone Dibati Worda Modermo, Bechati, Legbuna, Gireze and Zegeh Kebele (Bari Villige Peoples) for their providing the necessary information about use of medicinal plant on the study area during data collection. The people of Dibati District including the local authorities were also acknowledged for their positive response, sharing their valuable knowledge and times as well as for their tremendous generosity and hospitality. Dr. Tena Regassa deserved our heartfelt gratitude for their help in identification of plant specimens. Finally, our sincere appreciation goes to Mr. Gadissa who prepared the local map of the study District.

References

- [1] Abiyu Enyew, Zemed A, Ensermu K, Raja N (2014). Ethnobotanical Study of Traditional Medicinal Plants in and Around Fiche District, Central Ethiopia. *Current Research Journal of Biological Sciences* 6 (4): 154-167. Addis Ababa, Ethiopia, Uppsala Sweden 2009; 8: 1-7.
- [2] Assegid Assefa and Tesfaye Abebe. (2014). Ethnobotanical Study of Wild Medicinal Trees and Shrubs in Benna Tsemay District, Southern Ethiopia. *Journal of Science & Development* 2 (1) 17.
- [3] Ayeni. E and Basiri B. (2018). Ethnoveterinary Survey of Plants used in Treating Livestock among the Fulani people of Girei, Adamawa State, Nigeria. *WNOFNS*: 53-66 EISSN 2543-5426.
- [4] Balcha Abera (2014). Medicinal plants used in traditional medicine by Oromo people, Ghimbi District, Southwest Ethiopia. *Journal of Ethnobiology and Ethnomedicine*; 10 (40): 1-15.
- [5] Bayafers Tamene (2000). A Floristic Analysis and Ethnobotanical Study of the Semi- Wet land of Cheffa Area, South Wello, Ethiopia. *Agriculture and Healthcare*: ISSN 2224-3208 (Paper) ISSN 2225-093 Vol. 5, No. 1.
- [6] Bizuneh Woldeab, Reta Regassa, Tibebe Alemu, and Moa Megersa (2018). Medicinal Plants Used for Treatment of Diarrhoeal Related Diseases in Ethiopia. *Evidence-Based Complementary and Alternative Medicine Volume Article ID4630371*, 20 pages <https://doi.org/10.1155/2018/4630371>
- [7] Cotton, C. M., 1996. *Ethnobotany: Principles and Applications*. John Wiley and Sons LTD. Chichester, UK, pp. 424.
- [8] Endalew Amenu (2007). *Use and Management of Medicinal Plants by indigenous People of Ejaji Area (Chelya Wereda) West Shewa, Ethiopia: An Ethnobotanical Approach*, M.Sc Thesis. Addis Ababa Ethiopia.
- [9] Endeshaw Bekele (2007): study on actual situation of mechanical plants in Ethiopia prepared for Japan association for international collaboration of agriculturature and forestry.
- [10] Debela Hunde. (2001). *Use and Management of Traditional Medicinal Plants by Indigenous People of Bosat Woreda, Wolenchiti area: An ethnobotanical approach*. M.Sc. Thesis, Addis Ababa, Ethiopia.
- [11] Fisseha Mesfin (2007). *An Ethnobotanical Study of Medicinal Plants in Wonago Wereda, SNNPR, Ethiopia*, M.Sc. Thesis. Addis Ababa University, Addis Ababa. Ethiopia.
- [12] Friedman, J. Yaniu, Z., Dafni, A. and Palewitch, D. (1986). A Preliminary Classification of the Haling Potential of Medicinal Plants, Based on the Rational Analysis of Ethnopharmacolgical Survey Among Bedouins in Negue Desert, Israel. *Journal of Ethnopharmacology* 16: 275-2.
- [13] Getnet Chokole, Zemed A, Ensermu K (2015). Ethnobotanical study of medicinal plants in the environs of Tara-gedam and Amba remnant forests of Libo Kemkem District, northwest Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 2015; 11 (4): 1-38.

- [14] Getnet Chekol (2017). Ethnobotanical study of medicinal plants used against human ailments in Gubalafto District, Northern Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 13: 55.
- [15] Gidey Yirga (2010). Use of traditional medicinal plants by indigenous people in Mekele town, capital city of Tigray regional state of Ethiopia, *Journal of Medicinal Plants Research* 4: 25-50.
- [16] Kebu Balemie, Ensermu Kelbessa and Zemede Asfaw., (2004). Indigenous medicinal plant utilization, management and threats in Fentalle area, Eastern Shewa, Ethiopia. *Ethiopian Journal Biological Science*, 3: 37-58.
- [17] Luiz. R. Saldanha. G., Reinaldo, F., Paiva deA. (2005). Knowledge and use of medicinal plants by local specialist in a region of Atlantic Forest in the state of Pernambuco (North eastern Brazil).
- [18] Martine G (1995). *Ethnobotany: A method of manual* Chopman and Hall London UK. Grat Britain University Press e Cambridg press p 267. May 1998, IBCR, Addis Ababa, Ethiopia.
- [19] Mirutse Giday, Tilahun Teklehaymanot, Abebe Animut, and Yalemtehay Mekonn (2007). Medicinal plants of the Shinasha, Agew-awi and Amhara peoples in northwest Ethiopia. *Journal of Ethnopharmacology* 110: 516–525.
- [20] Moa Megersa (2010) *Ethnobotanical Study of Medicinal Plants in Wayu Tuka Wereda, East Wollega Zone of Oromia Region.*, MSC Thesis Addis Ababa, Ethiopia.
- [21] Mulugeta Kebebew (2016). Knowledge of medicinal plants used in and around Fincha'a Town, Western Ethiopia. *Journal of Pharmacognosy and Phytochemistry*; 5 (6): 110-114.
- [22] Nemarundwe, N. and Richards, M. (2002). Participatory methods for exploring livelihood Values derived from forests: potential and limitations. London. Pp. 168-198.
- [23] Nigatu Tuasha, Beyene Petros and Zemede Asfaw (2018). Medicinal plants used by traditional healers to treat malignancies and other human ailments in Dalle District, Sidama Zone, Ethiopia. *Journal of Ethnobiology and Ethnomedicine*: 14: 15-22.
- [24] Nigussie Amsalu, Yilkal Bezie, Mulugeta Fentahun, Addisu Alemayehu, and Gashaw Amsalu (2018). Use and Conservation of Medicinal Plants by Indigenous People of Gozamin Wereda, East Gojjam Zone of Amhara Region, Ethiopia: Article ID 2973513, 23 pages <https://doi.org/10.1155/2018/2973513>
- [25] Njau, A. E., 2001. An Ethnobotanical study of Medicinal Plants used by the Maasai People of Manayara, Arusha, Tanzania. M.Sc. Thesis. Addis Ababa, Ethiopia.
- [26] Tadesse Birhanu, Dereje Abera and Eyasu Ejeta (2015). Ethnobotanical Study of Medicinal Plants in Selected Horro Gudurru Woredas, Western Ethiopia., *Journal of Biology, Agriculture and Healthcare*: ISSN 2224-3208 (Paper) ISSN 2225-093X (Online) Vol. 5, No. 1.
- [27] Tafesse Mesfine and Mekonnen Lemma (2001), the role of traditional veterinary Herbal medicinal and its constraints in the animal health care system in Ethiopia, Addis Ababa Ethiopia.
- [28] Tena Regasa, (2015). Vascular plant diversity and ethnobotanical study of medicinal and wild edible plants in Jibat, Gedo and Chilimo forests, West Shewa Zone of Oromia Region, Ethiopia, PhD Dissertation, Addis Ababa, Ethiopia.
- [29] Tolosssa. k, Debela. E Athanasiadous. S, Tolora. A., Ganga. G, Houdijk, J (2013) ethno-medicinal study of plant used for treatement of human and livestock aliment by traditional healer in south omo, southern Ethiopia. *Ethio biol. ehtnomed*. 9: 32.
- [30] Yamane T (1967) *Statics, an Introduction Analysis*, 2nd Ed., New York: Harper and Row.
- [31] Yihenew Simegniew Birhan, Sintayehu Leshe Kitaw, Yihalem Abebe Alemayehu, Nakachew Minuye Mengesha (2018), Ethnobotanical study of medicinal plants used to treat human diseases in Enarj Enawga District, East Gojjam Zone, Amhara Region, Ethiopia. *SM J Med Plant Stud.*; 2 (1): 1006.
- [32] Zerhiun W, Mesfin T (1990). The Status of the Vegetation in the Lake region of the Rift Valley of Ethiopia and Possibilities of its Recovery. *SINET: Eth. J. Sci.*, 392: 97-120.
- [33] Zemede Asfaw (2001). The role of home gardens in production and conservation of medicinal plants. In: *Conservation and Sustainable Use of Medicinal Plants in Ethiopia, Proceeding of The National Work Shop on Biodiversity and Sustainable use of Medicinal Plants In Ethiopia*, 2.